Hypothyroidism

Hypothyroidism results from under-secretion of the hormone from the thyroid gland. The commonest cause globally, is primary thyroid failure due to chronic autoimmune thyroiditis (Hashimoto thyroiditis). Other well known causes are surgical removal, radioactive iodine therapy and external irradiation of the gland. Lithium and Interferon are drugs known to cause hypothyroidism. Secondary thyroid failure, from hypothalamic or pituitary disease, is rare. Patients should undergo assessment to determine the cause of hypothyroidism.

Clinical features

The symptoms depend on the duration and severity of hypothyroidism, and the rapidity with which the deficiency occurs. The psychological state of the patient too contributes to the clinical picture.

Box 1. Clinical features of hypothyroidism

Symptoms	Signs
Tiredness	Goitre
Weight gain	Bradycardia
Cold intolerance	Delayed relaxation of deep tendon reflexes (hung-up reflex)
Somnolence	Oedema
Dry skin	Slow speech
Coarseness and loss of hair	Dry skin
Hoarseness	Pallor
Constipation	Yellow skin
Impairment of memory	Ataxia
Myalgia and cramps	
Depression	
Menstrual disturbances in women	

Patients with chronic thyroiditis have a high incidence of other associated autoimmune diseases such as vitiligo, rheumatoid arthritis, Addison disease, diabetes mellitus, and pernicious anaemia.

Instances where referral to an endocrinologist is advisable include hypothyroidism in children, patients not responding to therapy or with structural changes in the thyroid gland, pregnancy, patients with cardiac disease and presence of other endocrine disease.

Laboratory diagnosis

The most cost-effective test is a sensitive measurement of TSH (thyroid stimulating hormone) level. A free T_4 (FT_4) level helps to confirm the diagnosis. An elevated

TSH and a decreased FT_4 are the expected findings. When autoimmune thyroiditis is suspected either antimicrosomal or antithyroglobulin antibody levels are done for confirmation. Rarely, patients with chronic thyroiditis have a change from a hypothyroid to a non-suppressible euthyroid state, or even to a hyperthyroid state, because of the development of stimulating TSH receptor autoantibodies (TSI or TRAb) of Graves disease.

Other biochemical abnormalities in hypothyroidism include elevated serum cholesterol, creatine phosphokinase and lactate dehydrogenase. Electrocardiographic changes are bradycardia, low amplitude QRS complexes and flattened or inverted T waves.

Treatment

Clinical hypothyroidism is treated with levothyroxine. Preferably, the patient should receive the same brand of levothyroxine throughout treatment. The mean daily replacement dose is 1.6 $\mu\text{g}/\text{kg}$ body weight, for an otherwise healthy adult under 50 years of age. The appropriate pace of treatment depends on the duration and severity of the hypothyroidism, and on the presence of other medical disorders. The dose should be gradually increased till full replacement is achieved.

In patients under 50 years who are free of other medical disorders full replacement dose is given in the first instance. Children may require higher doses (up to 4 $\mu\text{g}/\text{kg}$ body weight). In patients above 50 years of age or with cardiac disease, a lower initial dose is indicated (12.5-50 $\mu\text{g}/\text{day}$). Currently, there is insufficient evidence as to which patients with hypothyroidism will be better treated with T_3/T_4 combination than T_4 alone.

Follow up

Follow up includes assessment of response to therapy, drug interactions and adjustment of dose according to changes in body weight. Initial assessment should take place 6 weeks after starting therapy. Serum TSH is the most important investigation, and a FT_4 estimate may be included in the assessment. Dose adjustments should be carried out at 4-6 weekly intervals, with dose increments of 25-50 μg until TSH becomes normal. Once TSH is normal, the patient can be seen at 6-12 monthly intervals. Thyroxine replacement should be lifelong.

Special considerations

Subclinical hypothyroidism

This is defined as the presence of an elevated serum TSH concurrently with a normal FT_4 . Some patients with this combination feel clinically better when treated with levothyroxine. Treatment is advisable, especially if the thyroid antibodies are positive. If they are not treated,

frequent follow up should be done to detect the development of overt hypothyroidism.

Myxoedema coma

This is a life-threatening condition, which is generally seen in ill, elderly people with long-standing untreated hypothyroidism. Clinical features include coma, hypothermia, bradycardia, and respiratory and cardiac failure. Treatment is with intravenous levothyroxine and steroids, preferably in an intensive care unit.

Chronic thyroiditis with stimulating thyroid antibodies

Rarely, patients with chronic thyroiditis have a

change from a hypothyroid to non-suppressible euthyroid state or even to a hyperthyroid state because of the development of stimulating TSH receptor autoantibodies (TSI or TRAb) of Graves disease. If such patients had been receiving levothyroxine treatment, dose reduction or even cessation of levothyroxine therapy might be required.

Elderly

Hypothyroidism is highly prevalent in women over 60 years. Therefore screening with serum TSH is recommended for them. Symptoms may be subtle in the elderly. The differential diagnoses include depression, Parkinson disease and Alzheimer disease.

Dinithi Fernando, MD (Colombo), MRCP (UK). *Senior Lecturer in Physiology, Faculty of Medicine, Colombo 8.*
e-mail: <gimhanafdo@yahoo.com> *Competing interests: none declared.*