Solitary nodule of the thyroid

Solitary nodule of the thyroid is defined as a palpably discrete swelling within an otherwise apparently normal thyroid gland. When a clinician finds such a nodule, there are four main questions to answer, though not necessarily in the following order. The management depends on the answers.

1. Is it really a solitary nodule or a dominant nodule of a multinodular goitre?
2. What is the thyroid status of the patient?
3. Is nodule benign or malignant?
4. Are there any other patient characteristics and preferences that may influence the management?

The clinical evaluation and the biochemical, histological and radiological information should provide the answers to above questions.

Clinical evaluation

The clinical history and physical examination alone are not reliable for diagnosis or excluding a malignancy. Physical examination should include assessment of the thyroid status, examination for regional lymphadenopathy and features of local invasion, and compression.

Box 1. When to suspect a cancerous nodule

Nodule occurring in

- extremes of age (<20 years and >60 years)
- men – particularly elderly
- pregnancy
- the presence of features of local invasion (hoarseness, dysphagia, shortness of breath)
- an individual exposed to external irradiation of head, neck and chest
- an individual with a family history of thyroid cancer or multiple endocrine neoplasia or Gardner syndrome

Box 2. Features indicating a benign lesion

- A sudden or gradual onset of pain or tenderness
- Symptoms of hypothyroidism
- Family history of benign thyroid nodular disease, Hashimoto thyroiditis

Figure. Management and follow up of a clinically solitary thyroid nodule

Physical characteristics of a nodule do not reliably indicate its histological nature. For example, a hard nodule can be due to chronic thyroiditis or calcified adenoma and a cystic papillary cancer may feel soft.

Laboratory examination

Measurement of the serum thyroid stimulating hormone concentration is necessary and usually the only blood investigation required to identify patients with unsuspected thyrotoxicosis. Serum calcitonin should be measured when medullary thyroid carcinoma is suspected.

Fine needle aspiration cytology (FNAC) has become the initial test in most patients with a solitary nodule or dominant nodule within a multinodular goitre. It is safe, inexpensive and provides a better selection of patients for surgery. With the use of FNAC, the number of patients undergoing surgery for solitary nodule has been halved. In those who do have surgery, twice as many have been found to have thyroid cancer. Diagnostic accuracy of FNAC is between 70-95%, provided it is performed and interpreted by experienced specialists.

Sampling errors occur in very small or very large nodules, haemorrhagic nodules or multinodular glands. Taking multiple specimens or using ultrasound guided biopsy may minimise this problem. Three cytological results are possible when a satisfactory aspirate is obtained: benign, malignant and indeterminate (suspicious). When the biopsy specimen is inadequate, repeat FNAC is recommended. (See leading article page 4-6).

Imaging

Radioisotope scans (iodine or technetium)

Radioisotope scans are unnecessary in most patients and should not be included in the routine evaluation. They cannot reliably distinguish thyroid cancer from non-cancerous nodules. However, they are useful in patients with indeterminate cytology: for example, if a suspicious nodule is hyperfunctioning, it is almost always benign, whereas a hypofunctioning nodule in a patient with Graves disease is likely to be malignant.

Ultrasound scan

This is a useful test to determine whether a nodule is solid, cystic or mixed, but it does not differentiate between benign and malignant lesions. It is useful when FNAC is technically difficult (complex cysts, difficultly palpable nodules, nodules detected incidentally during other imaging procedures)

Management

A benign nodule (negative cytology) can be safely followed up. Either observation or thyroxine therapy is advised, depending on patient characteristics and preferences. For example, long term thyroxine therapy can cause osteoporosis in post-menopausal women. The goal of thyroxine therapy is to reduce the size of the nodule and prevent further growth.

About 10% of cytological diagnoses are malignant (positive cytology) and need surgery.

When a specimen is suspicious or indeterminate, there is no effective method to diagnose cancer accurately and surgery is advised in most instances as histology will reveal a definitive diagnosis. Radionuclide scanning is useful to determine the functional status of a nodule.

Conclusions

- FNAC is the first line investigation in a euthyroid patient with a solitary nodule.
- Radionuclide imaging should be reserved for nodules with indeterminate cytology or associated with thyrotoxicosis.
- Surgery is indicated for malignant and indeterminate lesions.
- Benign nodules are treated with thyroxine or observation. Selection of patients depends on patient characteristics and preferences.

Dinithi Fernando, MD (Colombo), MRCP (UK). Senior Lecturer in Physiology, Faculty of Medicine, Colombo 8.
e-mail: <gimhanafdo@yahoo.com>. Competing interests: none declared.